


AD-A174 771 PLASMA OPIOID PEPTIDE RESPONSES DURING HEAT ACCLIMATION 1/1
IN HUMANS(U) ARMY RESEARCH INST OF ENVIRONMENTAL
MEDICINE NATICK MA W J KRAEMER ET AL 27 OCT 86
UNCLASSIFIED USARTEM-M-7-87 F/G 6/15 NL



Plasma Opioid Peptide Responses During
Heat Acclimation in Humans

AD-A174 771

William J. Kraemer¹, Lawrence E. Armstrong², Louis J. Marchitelli¹,
Roger W. Hubbard² and Natalie Leva²

Exercise Physiology Division¹ and Heat Research Division²
US Army Research Institute of Environmental Medicine
Natick, MA 01760-5007 USA

Running head: Endogenous opioid peptides and heat

Send all correspondence to:

William J. Kraemer, Ph.D.
Biochemistry Laboratory
Exercise Physiology Division
US Army Research Institute of
Environmental Medicine
Natick, MA 01760-5007

27 OCTOBER 1986

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

DTIC
ELECTE
DEC 5 1986

B

86 12 05 04

Ref
Kraemer, W.J., L.E. Armstrong, L. Marchitelli, R.W. Hubbard, and N. Leva. *Ref*
Plasma opioid peptide responses during heat acclimation. PEPTIDES.....Plasma β
endorphin, Met-enkephalin and Peptide F immunoreactivity (ir) were measured at
rest and following exercise on three days (~~days 1,4,8~~) of an eight day heat
acclimation regime. Fourteen male subjects demonstrated physiological heat
acclimation adaptations. Our data demonstrated a differential response of
peripheral plasma levels of endogenous opioid peptides (EOP) to exercise in the
heat. In addition, EOP did not follow the same time-course of other
physiological adaptations as no differences (~~day 1 vs 4 vs 8~~) in resting or exercise
levels were observed over the eight day heat acclimation regime. Significant
increases in β -endorphin ir (pre- to post-exercise) appear to reflect concomitant
exercise-heat related changes. Furthermore, the increased peripheral levels of β -
endorphin may be related to increased glucocorticoid activity. Heat and exercise
stress may result in a reduction of Met-enkephalin ir observed in peripheral
plasma and might be due to degradation or a decrease in processing from the
larger precursors. The differential responses of EOP suggest the possibility of
separate physiological roles for these peptides during exercise in the heat but
peripheral plasma levels of EOP do not appear to reflect heat acclimation
changes. *X*



β -endorphin, Met-enkephalin, Peptide F, cortisol, heat acclimation

Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
PER FORM 50	
By	
Distribution	
Availability	
Dist	Special
A-1	

Endogenous opioid peptides (EOP) appear to be involved with thermoregulation and adaptation to extremes in the thermal environment [2,14]. Studies using the opiate antagonist naloxone which examined hypothermic and hyperthermic actions, suggest that the enkephalins and β -endorphin may be working through separate opiate receptors [3,6,10,14]. Pharmacological studies have also supported possible roles for EOP in thermoregulation [7,24,25]. To date, most studies have shown EOP to be involved in central thermoregulatory effects, although the possibility of peripheral thermoregulatory effects have been considered (27). Increases in rectal temperature (T_{re}) from combined exercise and heat stress have been associated with increases in peripheral circulating plasma concentrations of β -endorphin/ β -lipotropin in humans (16). While these data all support a possible role for EOP in acute thermoregulation, no data are available concerning the responses of EOP in humans to repeated heat stress exposures.

Most heat acclimation adaptations, induced by exercise and artificial climatic conditions in an environmental chamber, have been shown to occur over a 3 to 8 day period (1). It can be achieved by exercising in moderate to hot ambient conditions for 1 to 2 hours daily (21). The commonly observed physiological adaptations to heat stress over repeated exposures include, a decreased exercising core temperature, a decreased exercising heart rate and a plasma volume expansion (1).

It has been firmly established that exercise can significantly perturbate thermoregulatory mechanisms. In addition, exercise previously has been demonstrated to increase various EOP levels such as Methionine-enkephalin (Met-enkephalin), Peptide F and β -endorphin [4,5,9,11,12,15,19]. It has been suggested that training may also effect exercise responses of β -endorphin and Peptide F

[4,19]. The purpose of this study was to examine the resting and exercise plasma responses of β -endorphin, Met-enkephalin, and Peptide F (preproenkephalin 107-140; 33 amino acids) to an eight day heat acclimation regime in order to: 1) determine if any changes in resting or exercise EOP levels occur with heat acclimation 2) document any differential responses of different EOP and 3) gain insights into possible mechanisms which might help explain plasma levels of EOP observed.

METHODS

Fourteen healthy male subjects volunteered as subjects for this study. The physical characteristics of the subjects were $\bar{x} \pm SD$: age (yrs) 28.4 ± 1.9 , height 177.0 ± 2.0 (cm), weight 79.77 ± 3.78 (kg), % body fat 18.7 ± 1.4 , maximal oxygen consumption 45.74 ± 1.96 (ml \cdot kg \cdot min $^{-1}$) and surface area 1.96 ± 0.50 (m 2). After informed consent was obtained, all subjects completed a health questionnaire, activity questionnaire and history of heat exposure prior to testing. Data from these forms were examined to determine if each subject was unacclimatized prior to the study.

One day prior to the eight day heat acclimation regime, each subject was given a progressive exercise test to determine maximum oxygen consumption ($\dot{V}O_{2max}$), which was used to calculate the relative exercise intensities of the exercise training. During testing and training a semi-automated system was used to collect and analyze expired gases. This system consisted of a Hewlett-Packard 85B computer, scanner, and digital voltmeter were interfaced with a gas meter (Parkinson-Cowan), oxygen analyzer (Applied Electrochemistry S3A) and carbon dioxide analyzer (Beckman LB2). Tre was monitored every four minutes using a rectal probe (inserted 8 cm beyond the anal sphincter).

The heat acclimation regime consisted of eight days of treadmill exercise (0% grade). Each exposure consisted of 50 minutes of intermittent exercise during 100 minutes in an environmental chamber maintained at $41.2 \pm 0.5^{\circ}\text{C}$, $39.0 \pm 1.7\%$ relative humidity (RH). Exercise protocols on days 1 and 8 were performed at the identical duration and intensity ($71.8\% \pm 2.9 \dot{V}\text{O}_{2\text{max}}$ for running). Day 4 running exercise was performed at the intensity of ($67.6\% \pm 2.3 \dot{V}\text{O}_{2\text{max}}$). The other days consisted of similar high intensity interval exercise. Water was drunk ad libitum throughout all trials, but could not be sprayed or poured on the body.

Subjects were encouraged to drink adequate water when they were not in the climatic chamber. Upon arriving at the testing site each day, subjects were weighed and produced a urine sample which was analyzed for specific gravity. If any subject had a urine specific gravity greater than 1.030, he drank water until a more dilute urine was produced (specific gravity < 1.030).

A 20 minute standing equilibration period in the heat preceded each antecubital blood sample (days 1, 4 and 8). A second antecubital blood sample was taken immediately post-exercise. Blood samples for measurement of plasma cortisol, Met-enkephalin and β -endorphin were collected into specialized chilled glass vacutainers containing the anticoagulant EDTA (7.2 mg/5ml whole blood). Blood was mixed gently and immediately centrifuged for fifteen minutes at $760 \times g$, 4°C . Blood samples for measurement of plasma Peptide F were collected in chilled glass vacutainers containing sodium heparin and $25 \mu\text{l/ml}$ whole blood of aprotinin (Sigma Chemical Co., St. Louis, MO), gently mixed and centrifuged at $1500 \times g$, 4°C for fifteen minutes. Plasma samples were stored at -115°C until analyzed. Hemoglobin was analyzed using the cyanmethemoglobin method (Hycel Inc., Houston, TX) and hematocrit was analyzed in triplicate using a micro-capillary technique. Serum sodium (Na^{+}) and potassium (K^{+}) were determined

using a flame photometer (Rainin Instruments, FLM3) . Changes in plasma volume (% Δ VPV) were calculated from changes in hematocrit and hemoglobin (8).

Ten μ l samples were used in duplicate to perform the radioimmunoassay (New England Nuclear, North Billrica, MA) for plasma cortisol immunoreactivity. Five hundred μ l of tracer and antiserum complex were added to the sample tubes simultaneously by use of a Beckman Accu-prep 221 automatic dilutor. Tubes were vortexed and incubated at room temperature for 30 minutes. The tubes were then centrifuged at 1500 x g for 10 minutes at 4° C, the supernatant decanted and pellets were counted for radioactivity.

Prior to radioimmunoassay (Immuno Nuclear Corp., Stillwater, MN), Met-enkephalin was extracted from the plasma using ODS-silica columns. One ml of plasma was acidified with 100 μ l of 1 M HCl and loaded on the column. The Met-enkephalin was then eluted off the column with a total of 4 mls of methanol. The methanol eluate was evaporated to dryness at 37° C. Samples were reconstituted with 1 ml of BSA-phosphate. Samples were vortexed and placed at 37° C for 10 min followed by another vortex. 200 μ l aliquats were assayed in duplicate. The radioimmunoassay employed simultaneous addition of sample, rabbit anti-met-enkephalin antibody, and ¹²⁵I met-enkephalin followed by an overnight incubation at 4° C. Cross-reactivity of met-enkephalin antibody was 2.8% with Leu-enkephalin and less than 0.002% with any of the other known opioid peptides.

Prior to radioimmunoassay (Immuno Nuclear Corp., Stillwater, MN), one ml of plasma was concentrated and β -endorphin was extracted using a column of β -endorphin antibody coated Sepharose particles. β -endorphin was eluted from the column with a total of 0.5 ml 0.025 N HCl and 200 μ l were immediately assayed in duplicate. The radioimmunoassay involved a disequilibrium method based on

an antibody with high sensitivity to β -endorphin. The sample and first antibody were incubated for 20 hours at 4° C. ^{125}I β -endorphin was added followed by a second incubation for 20 hours at 4° C. Phase separation was completed in 20 minutes with a pre-precipitated complex of second antibody, carrier and PEG added in a single pipetting step. Crossreactivity of the β -endorphin antibody was less than 5% with β -lipotropin and less than 0.01% with any other known endogenous opioid peptide.

In order to avoid non-specific displacement in the radioimmunoassay, the Peptide F from each sample was partially purified using HPLC type minicolumns. The methods used to purify the samples and conduct the radioimmunoassay were previously described [19,20]. The mean percent recovery of radioactively labeled Peptide F with this procedure was 86%. Peptide F was measured by radioimmunoassay in duplicate using commercially available ^{125}I ligand and antisera (Peninsula Laboratories, Belmont, CA). The antisera showed crossreactivities of less than 0.05% with any known opioid peptide. The plasma immunoreactivity showed parallel displacement to Peptide F.

All samples for each specific radioimmunoassay were measured in the same assay to avoid run to run assay variations. Determinations of the different plasma ir were accomplished with the use of a Beckman 5500 Gamma counter and data reduction system.

Statistical evaluation of the data was accomplished by using a 3 x 2 Analysis of Variance (ANOVA) (days x pre/post). Subsequent post hoc analysis was performed using a Tukey test. Additionally, Pearson product moment correlation coefficients were calculated for the data set. In this study, significance was chosen as $p < 0.05$.

RESULTS

All of the subjects were unacclimatized at the beginning of this investigation. No significant differences were found when comparing mean entering body weight or mean entering urine specific gravity across the eight days. No significant differences were observed between the running intensities chosen on the three different days.

To examine the physiological effects of daily heat acclimation trials, a number of physiological variables were compared. The following variables all demonstrated significant decreases in mean values from day 1 to day 8; final exercise heart rate (170 ± 3 vs 144 ± 5 bpm), Δ heart rate (rest to end of exercise (84 ± 3 vs 68 ± 6 bpm), final Tre (39.17 ± 0.10 vs 38.52 ± 0.16 °C) and Δ Tre (rest to end of exercise) (2.04 ± 0.09 vs 1.46 ± 0.16 °C). Resting, pre-exercise plasma volume changes and expansion were compared from days 1 through 8. Plasma volume significantly expanded (+5.9%) during the first four days of heat acclimation and stabilized through day 8 (+5.2%). There was also a significantly better defense of plasma volume during exercise on day 8 than on day 1 (-5.1 ± 1.1 vs $-7.1 \pm 0.9\%$).

No significant differences were found in serum sodium between day 1, 4 or 8 for pre-or post-exercise values (day 1, pre= 141 ± 1 , post= 140 ± 1 ; day 4, pre= 141 ± 1 , post= 140 ± 1 ; day 8, pre= 140 ± 1 , post= 141 ± 1 mmole/L). Serum potassium concentrations were significantly higher post-exercise on each of the three days, but no differences between days were observed (day 1, pre= 4.3 ± 0.1 , post= 4.7 ± 0.1 ; day 4, pre= 4.5 ± 0.1 , post= 4.7 ± 0.1 ; day 8, pre= 4.4 ± 0.1 , post= 4.8 ± 0.1 mmole/L).

There were significant increases in plasma cortisol in pre- to post- exercise on each day but there were no significant differences between days for resting or

exercise values (Figure 1). No significant differences were observed in Met-enkephalin ir and Peptide F ir (pre- to post-exercise in the heat) or between-days for resting or exercise values (Figures 2 and 3). β -endorphin significantly increased pre- to post-exercise on each day but no significant differences were observed between days (Figure 4). A significant correlation ($r=0.45$) was observed between all post-exercise cortisol and post-exercise β -endorphin concentrations. If β -endorphin values were grouped into different levels above rest (i.e. 2 fold increases, 3 fold increases etc.), the correlations between β -endorphin and cortisol continued to increase in each group up to $r=0.77$, $p<0.05$ for the 10 fold increase group. The data set produced no other significant intercorrelations.

DISCUSSION

Comparison of the physiological measurements on days 1 and 8 (identical trials) indicated that typical heat acclimation adaptations (decreased exercising core temperature, a decreased exercising heart rate and plasma volume expansion) had occurred (1). Despite these heat adaptations, no changes in resting or exercise responses of any EOP were observed over this period of time. In fact, the trend in the response was a tendency for the values to decline on day 8. Rather than speculate on a positive response during a longer heat acclimation period, the simplest explanation suggests that plasma levels of EOP do not respond to the acclimation process per se.

β -endorphin ir significantly increased pre to post-exercise on each of the three days examined. This is consistent with a response pattern keyed predominantly to exercise stress rather than thermoregulatory strain. Total heat stress is defined as the sum of the heat generated in the body (metabolic heat) plus the heat gained from the environment (environmental heat) minus the heat lost from the body to

the environment. The body's response to total heat stress is defined as heat strain. Since both environmental and metabolic heat stress remained constant and yet rectal temperature declined, the heat loss mechanisms improved to effect a reduction in heat strain. Thus, across time ambient and metabolic stressors remained constant but thermoregulatory strain declined. The response pattern of β -endorphin, pre- to post-exercise, is therefore consistent with the metabolic demands of the exercise intensity.

The post-exercise plasma concentrations of β -endorphin observed in this study were higher than previous values reported for similar exercise intensities in thermoneutral conditions [4,9,12,15]. The added thermal stress may have partially augmented the magnitude of the plasma concentrations observed. The increases observed in this study were even higher than a previous study examining β -endorphin/ β -lipotropin responses in humans to heat stress (16). Still, due to the higher thermal stress and exercise intensities used in the present study, our data appears to be consistent with these previous findings that peripheral plasma concentrations increase in response to increased exercise-thermal stress (16). The lack of any significant bivariate relationships to core temperature changes or exercise intensity suggests that the response may be due to a combination of factors including both heat and exercise stress.

An underlying mechanism that might help explain the higher concentrations in this study of β -endorphin in the peripheral plasma might be a change in the permeability of the blood-brain barrier. The blood-brain barrier has been hypothesized to retard peripheral plasma concentrations of β -endorphin (9). Recently, it has been hypothesized that the pituitary-adrenal axis may physiologically modulate the permeability of the brain microvasculature to

macromolecules (23). Furthermore, this may be related to such influences as circulating glucocorticoids. Although no cause and effect is implicated, our data showed a significant relationship between plasma cortisol and β -endorphin values. Furthermore, this relationship grew stronger as β -endorphin values increased above rest. Our data can not evaluate the mechanism(s) involved. Still, it would appear to support the need for further study in this area to determine the central and peripheral effects of glucocorticoids on the peripheral plasma levels of β -endorphin.

No significant differences were observed for Peptide F ir. Still, it is interesting to note that both resting and post-exercise values are higher than previously reported (19). No data exists examining Peptide F responses to heat stress. The adrenal medullary chromaffin cells have been shown to be a major source of enkephalin-containing polypeptides secreted by the same stimuli which induces epinephrine release in these same cells [17,22,26]. Previous work has demonstrated that passive heating (50°C) does not increase epinephrine values (18). Thus, higher resting levels of Peptide F ir observed may be an acute response to heat exposure and requires further study in relationship to epinephrine release.

Administration of met-enkephalin has been shown to effect body temperature (6). Its effects are related to the dose and hypothesized to be a function of central mechanisms [6,10]. The lack of any changes pre- to post-exercise in this study disagrees with previously reported increases of Met-enkephalin ir following exercise (15). This may be a function of degradation in the peripheral plasma (13). Still, the values reported in this study are much lower than other data examining exercise responses of Met-enkephalin in thermoneutral environments (15). The lower values observed in this study may be due to the higher concentrations of precursor proenkephalin fragments. This

possible mechanism is supported by the higher levels of Peptide F ir found both at rest and following exercise.

In summary, while the possible roles of EOP and thermal stress have been addressed before little data were previously available which have examined possible heat acclimation changes. Our data demonstrated that peripheral concentrations of different EOP do not reflect heat adaptations gained over an 8-day heat acclimation regime. Heat and exercise stress may result in a reduction of Met-enkephalin ir found in the peripheral plasma due to degradation or a decreased processing of larger precursor fragments. The different pattern of responses of β -endorphin compared to Met-enkephalin and Peptide F supports the possibility of separate physiological roles for these opioid peptides during exercise in the heat.

The views, opinions, and/or findings contained in this report are those of the authors and should not be construed as an official department of the Army position, policy or decision, unless so designated by other official documentation.

Human subjects participated in this study after giving their free and informed voluntary consent. Investigators adhered to AR 70-25 on Use of Volunteers in Research.

Acknowledgements: We would like to thank Dora Ward for her time in preparing the manuscript and Dr. Randolph V. Lewis, University of Wyoming for his helpful comments in the preparation of this manuscript.


REFERENCES


1. Armstrong, L.E. and J.E. Dziados, Effects of heat exposure on the exercising adult. In Sports Physical Therapy, edited by D.B. Bernhardt. New York: Churchill Livingstone, 1986, pp 197-215.
2. Brown, M.R. Thermoregulation. In: Brain Peptides, edited by D.T. Krieger, M.J. Browstein and J.B. Martin. New York: John Wiley and Sons, 1983, pp 302-312.
3. Bloom, A.S. and L.F. Tseng. Effects of beta-endorphin on body temperature in mice at different ambient temperatures. Peptides. 2(3) 293-297, 1981.
4. Carr, D.B, B.A. Bullen, G.S. Skrinar, M.A. Arnold, M. Rosenblatt, I.Z. Beitins, J.B. Martin, and J.W. McArthur. Physical conditioning facilitates the exercise-induced secretion of Beta-endorphin and Beta-lipotropin in women. New Engl. J. Med. 305: 560-563. 1981.
5. Colt, E.W.D., S.L. Wordlaw and A.G. Frantz. The effect of running on plasma- β -endorphin. Life Sci. 28: 1637-1640, 1981.
6. Clark, W.G. Emetic and hyperthermic effects of centrally injected methionine-enkephalin in cats. Proc. Soc. Exp. Biol. Med., 154: 540-542, 1977.
7. Clark, W.G. Influence of opioids on central thermoregulatory mechanisms. Pharmacol. Biochem. Behav. 10: 609-613, 1979.
8. Dill, D.B. and D.L. Costill. Calculation of percentage changes in volume of blood, plasma and red cells in dehydration. J. Appl. Physiol. 37: 247-248, 1974.
9. Farrell, P.A., W.K. Gates, M.B. Maksud, and W.P. Morgan. Increase in plasma β -endorphin/ β -lipotropin immunoreactivity after treadmill running in humans. J. Appl. Physiology: Respirat. Environ. Exercise Physiology. 52: 1245-1249, 1982.


10. Ferri, S., R. Arrigo Reina, A. Santagostino, G.M. Scoto and C. Spadaro. Effects of met-enkephalin on body temperature of normal and morphine-tolerant rats. Psychopharmacol. 58: 277-281, 1978.
11. Fraioli, F., C. Moretti, D. Paolucci, E. Alicicco, F. Crescenzi and G. Fortunio. Physical exercise stimulates marked concomitant release of β -endorphin and adrenocorticotrophic hormone (ACTH) in peripheral blood in man. Experientia, 36: 987-989.
12. Gamber, S.R., T.L. Garthwaite, C.H. Pontzer, E.E. Cook, F.E. Tristani, E.H. Duthie, D.R. Martinson, T.C. Hagen and D.J. McCarty. Running elevates plasma β -endorphin immunoreactivity and ACTH in untrained human subjects. Proc. Soc. Exper. Biol Med. 168: 1-4, 1981.
13. Goodman, R.R., L.D. Fricker, and S.H. Snyder, Enkephalins In: Brain Peptides, edited by D.T. Krieger, M.J. Brownstein and J.B. Martin. New York: John Wiley and Sons, 1983, pp 827-849.
14. Holaday, J.W., E. Wie, H.H. Loh and C.H. Li. Endorphins may function in heat adaptation. Proc. Natl. Acad. Sci. USA. 75: 2923-2927, 1978.
15. Howlett, T.A., S. Tomlin, L. Ngahfoong, L.H. Rees, B.A. Bullen, G.A. Skrinar and J.W. McArthur. Release of β -endorphin and met-enkephalin during exercise in normal women: response to training. British Medical Journal, 288: 1950-1952, 1984.
16. Kelso, T.B., W.G. Herbert, F.C. Gwazdauskas, F.L. Gross and J.L. Hess. Exercise-thermoregulatory stress and increased plasma β -endorphin/ β -lipotropin in humans. J. Appl. Physiol.: Respirat. Environ. Exercise Physiology, 57: 444-449, 1984.
17. Kilpatrick, D.L., R.V. Lewis, S. Stein and S. Udenfriend. Release of enkephalins and enkephalin-containing polypeptides from profused beef adrenal glands. Proc. Natl. Acad. Sci. USA. 77: 7473-7475, 1981.


18. Kolka, M.A. and L.A. Stephenson. Thermoregulation during active and passive heating during the menstrual cycle. The Physiologist, 28: 368, 1985.
19. Kraemer, W., B. Noble, K. Roberston and R.V. Lewis. Response of plasma proenkephalin peptide F to exercise. Peptides, 6, Suppl 2: pp 167-169, 1985.
20. Lewis, R.V., Enkephalin biosynthesis in the adrenal medulla. In: Advances in Biochemical Psychopharmacology, 33, edited by E. Costa and M. Trabucchi. New York: Raven Press, 1982, pp 167-174.
21. Lind, A.R. and D.E. Bass. Optimal exposure time for development of acclimatization to heat, Fed. Proc. 22: 704, 1963.
22. Livett, B.G., D.M. Dean, L.G. Whelan, S. Undenfriend and J. Rossier. Co-release of enkephalin and catecholamines from cultured adrenal chromaffin cells. Nature, 289: 317-318, 1981.
23. Long, J.B. and J.W. Holaday. Blood-drain barrier: Endogenous modulation by adrenal-cortical function. Science, 227: 1579-1583, 1985.
24. Nemeroff, C.B., A.J. Osbahr, P.J. Manberg, G.N. Ervin and A.J. Prange. Alterations in nociception and body temperature after intracisternal administration of neurotensin, β -endorphin, other endogenous peptides, and morphine. Pro. Natl. Acad. Sci. USA, 76: 5368-5371, 1979.
25. Viveros, O.H., E.J. Dilberto, E. Hazum and K.J. Chang. Opiate-like materials in the adrenal medulla: Evidence for storage and secretion with catecholamines. Mol Pharmacol, 16: 1101-1108, 1979.
26. Wong, T.M., A. Koo and C.H. Li. β -endorphin vasodilating effect on the microcirculatory system of hamster cheek pouch. Int. J. Peptide Protein Res. 18:420-422, 1981.

- Figure 1. Plasma cortisol immunoreactivity responses to eight days of heat acclimation. ▲ indicates a significant ($p < 0.05$) difference pre- to post-exercise.
- Figure 2. Plasma Met-enkephalin immunoreactivity responses to eight days of heat acclimation.
- Figure 3. Plasma Peptide F immunoreactivity responses to eight days of heat acclimation.
- Figure 4. Plasma β -endorphin immunoreactivity responses to eight days of heat acclimation. ▲ indicates a significant ($p < 0.05$) difference pre- to post-exercise.









END

1-87

DTIC